

Research article

Submitted: January 15th, 2025 – Accepted: April 25th, 2025 – Published: June 30th, 2025
DOI: 10.13133/2284-4880/1734

What's on the menu today? First report of nectarivory for *Rhynocoris cuspidatus* (Hemiptera: Reduviidae)

Maria PIZARRO-BORRULL¹, Mario ALAMO^{1, *}

¹ Entomology Section, Asociación Iberozaa, Madrid, Spain – mariapibo@gmail.com; ORCID: <https://orcid.org/0009-0009-5774-3503>;
mario.alamodelolmo@hotmail.com; ORCID: <https://orcid.org/0000-0002-7410-8511>

*corresponding author

Abstract

This study reports the first observation of nectarivory in the predator reduviid *Rhynocoris cuspidatus* (Ribaut, 1921) in Spain. One individual of *R. cuspidatus* was observed sucking nectar from a *Jacobaea vulgaris* Gaertn. inflorescence in a grassland meadow in Berrecil de la Sierra (Spain). Our observation suggested that *R. cuspidatus* can use floral resources to obtain sugar or moisture during extreme climate conditions, such as can occur during Mediterranean summer.

Keywords: Mediterranean ecosystems, Zoophytophagy, Assassin bug, Trophic interactions, Floral resources scarcity.

Introduction

The use of auxiliary trophic resources can be crucial for survival in climates such as the Mediterranean, where summers are characterized by a marked scarcity of trophic and water resource (García-Ruiz et al. 2011; Lionello et al. 2006; Seager et al. 2019). Under such extreme conditions, heteropterans are compelled to explore alternative food sources, demonstrating remarkable trophic plasticity. For instance, some species have been observed shifting their trophic roles, with herbivores feeding on carrion (Alamo & Cepeda 2024) or carnivores consuming plant-based resources, such as through phytophagy or nectarivory. These unconventional strategies have been shown to confer significant benefits, including increased survival, higher fecundity rates, and shorter development times (Coll 1998; Naranjo & Gibson, 1996; Ruberson et al. 1986).

One such adaptive strategy observed in predatory insects is zoophytophagy, where carnivores exploit plant-based resources under extreme conditions (Torres & Boyd 2009). These interactions can involve feeding on plant tissues (phytophagy) or consuming nectar from flowers (nectarivory). Reports of phytophagy include *Podisus maculiventris* nymphs (Pentatomidae) (Ruberson et al. 1986), *Geocoris punctipes* and *G. pallens* (Geocoridae) (York, 1944), and *Nesidiocoris tenuis* (Miridae) (Sanchez

2008). Nectarivory has also been observed in *Heniarthes erythromaeus* and *Zelus armillatus* (Reduviidae) (Gil-Santana & Keller 2022).

Reduviidae (Hemiptera: Heteroptera), also known as assassin bugs, is one of the most diverse groups of Heteroptera, with approximately 6,800 described species distributed worldwide (Hwang & Weirauch 2012), being the largest family of predaceous terrestrial Heteroptera (Costa et al. 2022).

Reduviids are obligate zoophages with remarkable predatory behaviour (Guillermo-Ferreira et al. 2012; Torres & Boyd 2009). Most reduviids exhibit a generalist diet, but some specialize in certain taxonomic groups (Evangelin et al. 2014). Nonetheless, cases of phytophagy and nectarivory have been observed, particularly among species in the Harpactorini tribe of the Harpactorinae subfamily (Gil-Santana & Keller 2022). *Atrachelus cinereus* (Fabricius, 1798) has been recorded feeding on pollen and plant seeds (Stoner et al. 1975), *Atopozelus opsimus* (Elkins, 1954) on extrafloral nectaries (Guillermo-Ferreira et al. 2012), and *Zelus versicolor* (Herrich-Schaeffer, 1848) sucking nectar from *Oxypetalum balansae* (Gil-Santana & Keller 2022), among others.

The genus *Rhynocoris* (Reduviidae: Harpactorinae) comprises circa 150 species worldwide (Putshkov & Moullet 2009), four of which are present in the Iberian Peninsula: *Rhynocoris annulatus* (Linnaeus, 1758), *R. cuspidatus*

Ribaut, 1921, *R. erythropus* (Linnaeus, 1767), and *R. iracundus* (Poda, 1761) (Goula et al. 2020; Vivas & López Gallego 2013). However, only one record of phytophagy (nectarivory) has been documented for this genus: *R. erythropus* feeding on *Verbascum* sp. flowers in southern Spain (Baena 2011).

This study presents the first observation of nectarivory in *Rhynocoris cuspidatus* (Ribaut, 1921) feeding on capitula of *Jacobaea vulgaris* Gaertn during the Mediterranean summer in the Iberian Peninsula, highlighting a zoophytophagous strategy in this carnivorous species.

Material and Methods

The study was conducted in the vicinity of Arroyo de la Angostura (Becerril de la Sierra, Madrid, Spain; 40.725917N, 3.9925W) (Fig. 1). Plant diversity was assessed during the last week of June 2023 using a 20 × 5 m transect, as part of a broader study on flower-but-



Fig. 1 – Habitat surrounding the recorded trophic interaction at Arroyo de la Angostura, 19th July 2023. © M-A

terfly trophic networks being carried out in the area of the interaction. The study site is characterized as open scrubland, dominated primarily by *Juniperus oxycedrus* and *Cistus ladanifer*, with less abundant representation of *Thymus vulgaris* and *Lavandula pedunculata*. The arboreal stratum is sparsely populated by holm oaks (*Quercus ilex*). By late June, the area exhibited a high diversity of floral resources, including both shrubs (*Rosa canina*, *Rubus ulmifolius*) and herbaceous species such as *Daphne gnidium*, *Leontodon saxatilis*, *Senecio squalidus*, *Anthemis arvensis*, *Crepis capillaris*, *Lavandula stoechas*, *Verbascum pulverulentum*, *Hirschfeldia incana*, and *Convolvulus arvensis*, among others. Plant species were identified using morphological keys (Flora Iberica) and geographic and visual identification systems (Pl@ntNet).

The streambed is not an active stream but rather an area with highly waterlogged soil. Human activities, primarily cattle grazing, are common in the study area.

The reduviid was identified through photographs taken in the field, following the taxonomic keys of Putshkov & Moulet (2009).

Results

The observation occurred at 10:43 a.m. on 19 July 2023, during a heat wave. Weather conditions were predominantly sunny, with 20–30% cloud cover, and a recorded temperature of 29.2°C (AEMET, 2025). An individual of *Rhynocoris cuspidatus* was observed feeding on an open capitulum of *Jacobaea vulgaris*. A significant amount of pollen was visible on the rostrum, indicating active feeding behavior (Fig. 2). Shortly after the photograph was taken, the individual ceased feeding and flew away.

Other *Jacobaea vulgaris* plants in the study area were found hosting *Tyria jacobaeae* caterpillars (Linnaeus, 1758) (Fig. 3). However, it is noteworthy that the *J. vulgaris* plant on which *Rhynocoris cuspidatus* was observed did not host any *T. jacobaeae* caterpillars. Most flowering plants recorded during June had either lost their flowers by the time of the observation, highlighting the scarcity of viable floral resources in the area during the peak of summer.

Discussion

Phytophagy and nectarivory behaviors have been reported in several reduviids, but there is limited information regarding the ecological context and motivations behind plant feeding in these generalist predators. Multiple hypotheses have been proposed, including the use of plants as additional sources of water or sugar (Baena 2011; Bérenger & Pluot-Sigwalt 1997; Stoner et al. 1975), as an




Fig. 2 – Observation of *Rhynocoris cuspidatus* with its rostrum covered in pollen after inserting it into the inflorescence of a capitulum of *Jacobaea vulgaris*. © M-A



Fig. 3 – *Jacobaea vulgaris* plant hosting *Tyria jacobaeae* caterpillars in the vicinity of Arroyo de la Angostura, 19th July 2023. © M-A

alternative resource during prey scarcity (Sanchez 2008), or as a safer feeding strategy compared to hunting (Guillermo-Ferreira et al. 2012).

Although *Rhynocoris cuspidatus* is primarily an insect predator, preying on species such as honey bees, its nectarivory behavior can be explained through several hypotheses. One possibility is that plant feeding provides an alternative source of water during arid conditions, as suggested for *R. erythropus* in Jaén, Spain (Baena 2011). However, despite the heat wave affecting the Iberian Peninsula, the study site maintained relatively moist conditions, likely due to water infiltration from Arroyo de la Angostura, which sustained a humid microenvironment. Given these conditions, it seems unlikely that the observed individual was feeding on *Jacobaea vulgaris*

solely to obtain water. Instead, feeding for sugar acquisition should be considered, as similar behaviors have been documented in other reduviids (Bérengrer & Pluot-Sigwalt 1997).

The observed individual of *Rhynocoris cuspidatus* was feeding on the capitulum of the toxic plant *Jacobaea vulgaris*, which contains pyrrolizidine alkaloids (PAs). These compounds are known to deter or intoxicate generalist herbivores (Harper & Wood 1954; Hartmann & Ober 2000; Kalač & Kaltner 2021). Despite its toxicity, some pollinators, such as Lepidoptera and Coleoptera, tolerate PAs and even utilize them as defense compounds against predators (Hartmann & Ober 2000; Nishida 2002). If *R. cuspidatus* was indeed using *J. vulgaris* as a food resource, it would likely require a tolerance mechanism for

PAs. However, the position of *R. cuspidatus* on the plant suggests that it may have been primarily using the capitulum as a hunting perch, with nectar feeding as a secondary behavior.

Many species of *Rhynocoris* (e.g., *R. iracundus*, *R. erythropus*), including *R. cuspidatus* (Baena 2011), have been observed using plant inflorescences as hunting spots, primarily preying on pollinators (Baena 2011; Gil-Santana & Alves 2011; Miller 1953; Putshkov & Moulet 2009). During this observation, other *Jacobaea vulgaris* plants in the study area hosted *Tyria jacobaeae* caterpillars, while the plant where *R. cuspidatus* was observed did not. This suggests that *R. cuspidatus* could potentially use *J. vulgaris* as a hunting platform to prey on *T. jacobaeae* caterpillars, which are the primary herbivores of this plant (Joshi & Vrieling 2005). If this were the case, *R. cuspidatus* would require a mechanism to tolerate PAs, as *T. jacobaeae* caterpillars sequester these toxic compounds from *J. vulgaris* (Aplin et al. 1968). However, no records of *R. cuspidatus* preying on these caterpillars currently exist.

By late July, flowering plants become scarce due to the summer drought and intensive grazing in the area. Under these conditions, non-palatable plants such as *Jacobaea vulgaris* and other Asteraceae were among the few floral resources available, limiting access for animals. In this context, *Rhynocoris cuspidatus* likely used *J. vulgaris* both as a secondary food resource for sugar and as a hunting platform, waiting for potential prey such as butterflies and beetles feeding on the plant's flowers.

This is the first report of *Rhynocoris cuspidatus*, a predatory reduviid, feeding on nectar from *Jacobaea vulgaris* in Spain. The evidence suggests that the reduviid utilized the plant as an alternative resource to obtain sugar and moisture during the Mediterranean summer, while simultaneously using the capitulum as a hunting perch.

Acknowledgements – This study was conducted during butterfly-flower interaction samplings, in collaboration with Hugo Alejandro Álvarez and Robert J. Wilson, which were funded by the project PID2021-126293OB-I00; MCIU/AEI/UE, 2022-2025 (DRAMA) and by an extension grant from the Juan de la Cierva fellowship FJC2021-046506-I (MCIN/AEI/10.13039/501100011033/NextGenerationEU/PRTR), Ministerio de Ciencia e Innovación, Spain. We extend our gratitude to both for their guidance and collaboration throughout the fieldwork, which made this study possible.

Conflict of interest disclosure, Ethics approval statement

None conflicts to our knowledge, no special approval required.

Contribution of authors

Maria Pizarro-Borrull drafted the initial manuscript. Mario Alamo collected the field data and provided critical insights into the ecological interpretation of the observations. Both authors collaboratively revised and refined the manuscript, contributing to the development of the final version.

References

- Affouard A., Joly A., Lombardo J., Champ J., Goeau H., Chouet M., Gresse H., Botella C., Bonnet P. 2023. Pl@ntNet automatically identified occurrences. v1.8. Pl@ntNet. Dataset/Occurrence. <https://ipt.plantnet.org/resource?r=queries&v=1.8>
- Agencia Estatal de Meteorología. 2025. Climatic data for the station “Madrid - Alpedrete” on 17 July 2023. Available online at: <https://opendata.aemet.es/centrodedescargas/productosAEMET>
- Alamo M., Cepeda J.C. 2024. When life gives you carcasses: first record of necrophagy by *Coriomeris hirticornis* on European rabbit (*Oryctolagus cuniculus*) (Hemiptera: Coreidae). *Fragmenta entomologica*, 56: 123–128. Doi: <https://doi.org/10.13133/2284-4880/1676>
- Aplin R.T., Benn M.H., Rothschild M. 1968. Poisonous Alkaloids in the Body Tissues of the Cinnabar Moth (*Callimorpha jacobaeae* L.). *Nature*, 219(5155): 747–748. Doi: <https://doi.org/10.1038/219747a0>
- Baena M. 2011. Unusual feeding habits in four Iberian Heteroptera (Hemiptera). *Boletín de la Sociedad Entomológica Aragonesa*, 48: 399–401.
- Béranger J.-M., Pluot-Sigwalt D. 1997. Relations privilégiées de certains Heteroptera Reduviidae prédateurs avec les végétaux. Premier cas connu d'un Harpactorinae phytophage. *Comptes Rendus de l'Académie des Sciences-Series III-Sciences de la Vie*, 320: 1007–1012.
- Castroviejo S. (ed.). 1986-2012. *Flora iberica* 1-8, 10-15, 17-18, 21. Real Jardín Botánico, CSIC, Madrid.
- Coll M. 1998. Living and Feeding on Plants in Predatory Heteroptera, in: Coll M., Ruberson J.R. (Eds), *Predatory Heteroptera: Their Ecology and Use in Biological Control*. SPIE. Doi: <https://doi.org/10.4182/YGQF2785.1998.89>
- Costa J., Chiquetto-Machado P.I., Gil-Santana H.R. 2022. Predation strategies of *Harpactor angulosus* (Lepeletier & Serville, 1825) (Hemiptera: Reduviidae) on *Cladomorphus phyllinus* Gray, 1835 (Phasmatodea: Phasmatidae) in captivity. *Revista Chilena de Entomología*, 48(3): 531–543. Doi: <https://doi.org/10.35249/rche.48.3.22.07>
- Evangelin G., Horne B., Muthupandi M., William S.J. 2014. Feeding Behaviour of the Predatory Reduviid, *Rhynocoris kumarii* (Hemiptera: Reduviidae). *International Journal of Life Science*, 3 (2): 64. Doi: <https://doi.org/10.5958/2319-1198.2014.01087.2>
- García-Ruiz J.M., López-Moreno J.I., Vicente-Serrano S. M., Lasanta-Martínez T., Beguería S. 2011. Mediterranean water resources in a global change scenario. *Earth-Science Reviews*, 105 (3-4): 121–139. Doi: <https://doi.org/10.1016/j.earscirev.2011.01.006>
- Gil-Santana H.R., Alves R.J.V. 2011. Association between *Zelus versicolor* (Herrich-Schäffer) (Hemiptera, Reduviidae, Harpactorinae) and *Bidens rubifolia* Kunth (Asterales, Asteraceae). *EntomoBrasilis*, 4(1): 30–32. Doi: <https://doi.org/10.12741/ebrazilis.v4i1.95>
- Gil-Santana H.R., Keller H.A. 2022. New records of associations between species of Reduviidae (Hemiptera: Heteroptera) and

- plants in Argentina. *Revista Chilena de Entomología*, 48(1): 55–63. Doi: <https://doi.org/10.35249/rche.48.1.22.04>
- Goula M., Osorio Álvarez V., Tomàs J., Roca-Cusachs M. 2020. Llista dels Heteròpters de Catalunya (Insecta, Hemiptera, Heteroptera). Versió 2. Publicacions del Centre de Recursos de Biodiversitat Animal, Universitat de Barcelona, 6: 2020, 46 pp. Doi: <https://hdl.handle.net/2445/169573>
- Guillermo-Ferreira R., Cardoso-Leite R., Gandolfo R. 2012. First observation of alternative food usage (extrafloral nectar) by the assassin bug *Atopozelus opsimus* (Hemiptera, Reduviidae). *Revista Brasileira de Entomologia*, 56(4): 489–491. Doi: <https://doi.org/10.1590/S0085-56262012000400014>
- Harper J.L., Wood W.A. 1954. Biological flora of the British Isles – *Senecio jacobaea* L. *Journal of Ecology* 45: 617–637. Doi: <https://doi.org/10.2307/2256946>
- Hartmann T., Ober D. 2000. Biosynthesis and Metabolism of Pyrrolizidine Alkaloids in Plants and Specialized Insect Herbivores, in: Leeper F.J., Vederas J.C. (Eds), Biosynthesis, Topics in Current Chemistry. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 207–243. Doi: https://doi.org/10.1007/3-540-48146-X_5
- Hwang W.S., Weirauch C. 2012. Evolutionary History of Assassin Bugs (Insecta: Hemiptera: Reduviidae): Insights from Divergence Dating and Ancestral State Reconstruction. *PLoS ONE* 7(9), e45523. Doi: <https://doi.org/10.1371/journal.pone.0045523>
- Joshi J., Vrieling K. 2005. The enemy release and EICA hypothesis revisited: incorporating the fundamental difference between specialist and generalist herbivores. *Ecology Letters*, 8(7): 704–714. Doi: <https://doi.org/10.1111/j.1461-0248.2005.00769.x>
- Kalač P., Kaltner F. 2021. Pyrrolizidine alkaloids of European *Senecio/Jacobaea* species in forage and their carry-over to milk: A review. *Animal Feed Science and Technology*, 280, 115062. Doi: <https://doi.org/10.1016/j.anifeeds.2021.115062>
- Lionello P., Malanotte-Rizzoli P., Boscolo R., Alpert P., Artale V., Li L., Luterbacher J., May W., Trigo R., Tsimplis M., Ulbrich U., Xoplaki E. 2006. The Mediterranean climate: An overview of the main characteristics and issues, in: *Developments in Earth and Environmental Sciences*. Elsevier, pp. 1–26. Doi: [https://doi.org/10.1016/S1571-9197\(06\)80003-0](https://doi.org/10.1016/S1571-9197(06)80003-0)
- Miller N.C.E. 1953. Notes on the biology of the Reduviidae of Southern Rhodesia. *The Transactions of the Zoological Society of London*, 27(2): 541–672. Doi: <https://doi.org/10.1111/j.1096-3642.1953.tb00233.x>
- Naranjo S.E., Gibson R.L. 1996. Phytophagy in predaceous Heteroptera: effects on life history and population dynamics, in: *Zoophytophagous Heteroptera: Implications for Life History and Integrated Pest Management*. Proceedings Thomas Say Publications in Entomology, pp. 57–93.
- Nishida R. 2002. Sequestration of Defensive Substances from Plants by Lepidoptera. *Annual Review of Entomology*, 47(1): 57–92. Doi: <https://doi.org/10.1146/annurev.ento.47.091201.145121>
- Putshkov P.V., Moulet P. 2009. Hémiptères reduviidae d'Europe occidentale. *Fédération française des Sociétés de sciences naturelles*.
- Ruberson J.R., Tauber M.J., Tauber C.A. 1986. Plant Feeding by *Podisus maculiventris* (Heteroptera: Pentatomidae): Effect on Survival, Development, and Preoviposition Period. *Environmental Entomology*, 15(4): 894–897. Doi: <https://doi.org/10.1093/ee/15.4.894>
- Sanchez J.A. 2008. Zoophytophagy in the plantbug *Nesidiocoris tenuis*. *Agri and Forest Entomology*, 10 (2): 75–80. Doi: <https://doi.org/10.1111/j.1461-9563.2007.00357.x>
- Seager R., Osborn T.J., Kushnir Y., Simpson I.R., Nakamura J., Liu H. 2019. Climate Variability and Change of Mediterranean-Type Climates. *Journal of Climate*, 32 (10): 2887–2915. Doi: <https://doi.org/10.1175/JCLI-D-18-0472.1>
- Stoner A., Metcalfe A.M., Weeks R.E. 1975. Plant feeding by Reduviidae, a predaceous family (Hemiptera). *Journal of the Kansas Entomological Society*, 48: 185–188.
- Torres J.B., Boyd D.W. 2009. Zoophytophagy in predatory Hemiptera. *Brazilian Archives of Biology and Technology*, 52 (2): 1199–1208. Doi: <https://doi.org/10.1590/S1516-89132009000500018>
- Vivas L. 2013. Listado de especies de heterópteros de la Península Ibérica en Biodiversidad Virtual. Datos extraídos del Banco Taxonómico-Faunístico Digital de los Invertebrados Ibéricos (BTFDII), pp. 20–28, in: *Especial Nº 2 - Septiembre de 2013. Heteroptera*. BV news, Revista de la Asociación Fotografía y Biodiversidad, Barcelona. Asociación Fotografía y Biodiversidad, BiodiversidadVirtual.org. 47 pp.
- York G.T. 1944. Food studies of *Geocoris* spp., predators of the beet leafhopper. *Journal of Economic Entomology*, 37(1): 25–29. Doi: <https://doi.org/10.1093/jee/37.1.25>

