



# ANNALI DI BOTANICA

*Ann. Bot. (Roma)*, 2012, 2: 67–78

Journal homepage: <http://annalidibotanica.uniroma1.it>



## SYNTAXONOMY OF SUBALPINE TALL-GRASS COMMUNITIES (*CALAMAGROSTIETALIA VILLOSAE*) IN THE UKRAINIAN DISTRICTS OF THE EASTERN CARPATHIANS

IAKUSHENKO D.<sup>1\*</sup>, BURLAKA M.<sup>2</sup>, CHORNEI I.<sup>3</sup>, KVAKOVSKA I.<sup>4</sup>, SOLOMAKHA V.<sup>5</sup> & TOKARYUK A.<sup>3</sup>

<sup>1</sup> University of Zielona Góra, Zielona Góra, Poland

<sup>2</sup> M.G. Kholodny Institute of Botany, Kyiv, Ukraine

<sup>3</sup> Yuriy Fedkovych Chernivtsi National University, Chernivtsi, Ukraine

<sup>4</sup> Uzhans'kyj National Nature Park, Velykyi Bereznyi, Zakarpattia region, Ukraine

<sup>5</sup> Taras Shevchenko National University, Kyiv, Ukraine

\* Corresponding author: Telephone: +48-68-3282557; e-mail: dmytrok@bigmir.net

(RECEIVED 21 APRIL 2012; RECEIVED IN REVISED FORM 30 MAY 2012; ACCEPTED 06 JUNE 2012)

**ABSTRACT** –This paper presents a phytosociological survey of subalpine tall-grass communities within the order *Calamagrostietalia villosae* (class *Mulgedio-Aconitetea*) in the Ukrainian districts of the Eastern Carpathians. This syntaxonomical revision includes relevés from the mountain ranges Beskydy, Gorgany, Chornohora, Svydovets', Marmarosh, Hryniava, and Chyvchyny. The analysis was performed using numerical classification methods. Five associations within three alliances were distinguished and concisely characterized. Typification and name inversion of the association *Hyperico alpigeni-Calamagrostietum villosae* Pawłowski et Walas 1949 *nom. invers. propos.* was performed.

KEYWORDS: SUBALPINE, TALL-GRASS, *CALAMAGROSTIETALIA VILLOSAE*, *MULGEDIO-ACONITETEA*, SYNTAXONOMY, CARPATHIANS, UKRAINE

## INTRODUCTION

The summit vegetation has been unevenly described in the different parts of the Carpathians. While for the Western Carpathians detailed surveys are available (Kliment & Valachovič, 2007; Kliment et al., 2010), data from the Eastern Carpathians seem to be still unstructured (Coldea, 1991; Malynovski & Kricsfalussy, 2002). Some local associations were described but contrasting classification schemes were developed for different parts of the Eastern Carpathians. See for Bukovské vrchi in Slovakia (Hadač et al., 1988), for Bieszczady in Poland (Winnicki, 1999), for the Ukrainian Carpathians (Malynovski et al., 1992; Malynovski & Kricsfalussy, 2000; 2002), and for the Romanian Eastern Carpathians (Coldea, 1991, Sanda et al., 1997; 2008). Subalpine tall-grass and tall-forbs communities of the class *Mulgedio-Aconitetea* has been studied in the Ukrainian Carpathians since the 1930s by Czech (Domin, 1930; Deyl,

1940), Polish (Swiderski & Szafran, 1931; Pawłowski & Walas, 1949), Slovak (Hadač et al., 1995), and Ukrainian researchers (Malynovski, 1980; Malynovski et al., 1992; Malynovski & Tsaryk, 1995; Malynovski & Kricsfalussy, 2000; 2002; Solomakha et al., 2004; Klimuk et al., 2006; Iakushenko, 2007). Nevertheless, these communities in the region were poorly documented so far. Moreover, data from the Ukrainian part of the Eastern Carpathians were unknown for a wide range of researches and only minimally included into large-scale reviews (cf. Michl et al., 2010). The syntaxonomical status of these communities remained ambiguous.

The aim of this paper is to describe the diversity of *Calamagrostietalia villosae* communities in the Ukrainian part of the Eastern Carpathians, and to revisit the nomenclature of already established syntaxa and to propose a new comprehensive syntaxonomical scheme for communities in this area.

## MATERIALS AND METHODS

The study area is a part of the Eastern Carpathians located within the borders of Ukraine (Fig. 1). It consists of low and gently sloping mountain ranges running in a northwest-southeast direction. The highest mountain group is the Chornohora with Hoverla as highest peak (2061 m a.s.l.). The other ridges with subalpine vegetation are Marmarosh (Mt Pip Ivan Marmaros'kyj, 1936 m), Svydovets' (Mt Velyka Blyznytsia, 1883 m), Gorgany (Mt Velyka Syvulia, 1836 m), Chyvchyny (Mt Chyvchyn, 1769 m), Borzhava (Mt Stij, 1677 m), Hryniava (Mt Baba-Liudova, 1590 m), and Eastern Beskydy (Mt Pikuj, 1405 m). The bedrock mostly consists of Carpathian flysch, except the volcanic Vyhorlat-Guta ridge and the crystalline Marmarosh mountain range. The climate is temperate continental with a cool and relatively moist (430–950 mm) vegetative season and a moderate dry (260–580 mm) winter with frequent thaws. A subalpine vegetation type ("polonyny") dominates in the Ukrainian Carpathians from 1200 to 1800 m a.s.l. (Malinovsky, 1980; 2003; Malinovsky & Kricsfalussy, 2000).



Figure 1. Location of the studying area within Europe.

The nomenclature of vascular plants follows "Flora Europaea" (Tutin et al., 1968 – 1980). Subspecies and varieties were merged at species level. However, for cryptogams (mosses and lichens), we used also entries that were identified only at the rank of genera.

The syntaxonomical revision was performed including original data, sampled 2005 to 2010 and data published between 1934 and 1955. The phytosociological relevé sampling followed the Braun-Blanquet approach (Westhoff & van der Maarel, 1973). A data set of 149 relevés was used for the analysis after removal of published relevés without information about mosses and lichens.

The relevés were stored in the TURBOVEG database (Hennekens & Schaminée, 2001) and classified by TWINSPAN protocol (Hill, 1979) using JUICE 6.5 software package (Tichý, 2002). Diagnostic species were chosen on the basis of fidelity measure (Chytrý et al., 2002) with corrections based on literature data. The threshold value for a species considered as diagnostic was set at a Phi coefficient (multiplied by 100) more than 30. Species that were recorded in at least 61 % of the relevés of an association were considered constant for it. The classification follows the results of previous syntaxonomical revisions of the class *Mulgedio-Aconitetea* within Europe (Michl et al., 2010) and Western Carpathians (Kočí, 2001; 2007; Kliment & Jarolímek, 2003; Matuszkiewicz, 2006; Kliment & Valachovič, 2007; Šibíková et al., 2008; Kliment et al., 2010). In the diagnoses of the communities, the following abbreviations were used: art. – article of the International Code of Phytocoenological Nomenclature (ICPN) (Weber et al., 2000), dom. – dominating species, opt. – optimal conditions, p. p. – pro parte (partially), syn. – synonym, syntax. syn. – syntaxonomical synonym, rel. – relevé. The diagnostic and constant species were arranged alphabetically.

## RESULTS

Five distinct groups of relevés were obtained and their syntaxonomical ranking has been assessed. Five associations have therefore been assigned to the three following alliances: *Festucion carpatica* (carbonatophilous chionophilous tall-grass communities), *Calamagrostion arundinaceae* (species-rich tall-grass communities of relatively warm and dry sites), and *Calamagrostion villosae* (mesophilous chionophilous tall-grass meadows on acid silicate bedrock).

### ***MULGEDIO-ACONITETEA* HADAČ ET KLIKA IN KLIKA ET HADAČ 1944**

High-altitude tall-herb and tall-grass vegetation.

#### ***Calamagrostietalia villosae* Pawłowski et al. 1928**

Tall-grass mesophilous acidophilous to basiphilous vegetation of subalpine meadows.

#### ***Festucion carpatica* Bělohlávková et Fišerová 1989**

Chionophilous tall-grass communities on moist carbonate soils in montane and subalpine belts.

#### ***Festucetum carpatica* Domin 1925**

Species-rich chionophilous tall-grass communities dominated by the Carpathian endemic *Festuca carpatica* of montane and

subalpine belts (1450–1780 m) on moist, well drained neutrophilous or slightly basiphilous skeleton-rich carbonate soils, on steep (from 10–20° to 40–55°) sheltered slopes, glens and stabilized calcareous screes.

**Syn.:** *Carduo kernerii-Festucetum carpaticae* (Pușcaru et al. 1956) Coldea (1986) 1990 (syntax. syn.), *Festucetum carpaticae* (Domin 1925) Pawłowski et Stecki 1926 (syntax. syn.).

**Incl.:** *Thymo-Festucetum amethystinae festucetosum carpaticae* Kricsfalussy et Malynovski 2000, p.p.

**Nomenclatural type:** Domin, 1925: 8, rel. 3, lectotypus (Unar et al., 1985: 39).

**Diagnostic taxa:** *Aquilegia nigricans*, *Bupleurum longifolium*, *Astrantia major*, *Cirsium erisithales*, *Cortusa matthioli*, *Euphrasia picta*, *Festuca carpatica* (dom., opt.), *Galium anisophyllum*, *Linum catharticum*, *Listera ovata*, *Lotus corniculatus*, *Primula elatior*, *Ranunculus oreophilus*, *Rhinanthus alpinus*, *Saxifraga paniculata*, *Scabiosa columbaria*, *Thymus pulcherrimus*, *Veronica urticifolia* (Table 1).

**Constant taxa:** *Astrantia major*, *Calamagrostis villosa*, *Cirsium erisithales*, *Euphrasia picta*, *Festuca carpatica*, *Gymnadenia conopsea*, *Primula elatior*.

**Distribution in the Ukrainian Carpathians:** Sydovets', Chyvchyny Mts (Chorny Dil ridge) (Fig. 2, a).

**References:** Domin, 1925, 1930; Deyl, 1940; Malynovski, 1980; Unar et al., 1985; Bělohlávková & Fišerová, 1989; Sanda et al., 1997; Malynovski & Kricsfalussy, 2000; Kliment et al., 2007, 2010; Šibíková et al., 2008; Ustymenko & Tasenkevich, 2009; Michl et al., 2010.

### *Calamagrostion arundinaceae* (Luquet 1926) Oberdorfer 1957

Species-rich moderate thermophilous tall-grass communities in relatively warm and dry habitats in upper montane and subalpine belts.

**Syn.:** *Calamagrostion atlanticum* Luquet 1926 (art. 34a), *Calamagrostion arundinaceae* Oberdorfer 1949 (art. 8), *Calamagrostion arundinaceae* Jeník 1959 (art. 8), *Calamagrostion arundinaceae* (Luquet 1926) Jeník 1961 (art. 31).

### *Achilleo strictae-Calamagrostietum arundinaceae* Hadač et al. 1988

Species-rich tall-grass pasturelands (“polonya”) in montane and subalpine belts (850–1200 m) on gentle, relatively warm

semi-dry (southern, southeastern, and western) slopes on subacidic soils. These communities are usually dominated by *Calamagrostis arundinacea*, associated with Eastern-Carpathian and South-Eastern-Carpathian species.

**Syn.:** *Tanaceto-Calamagrostietum arundinaceae* Winnicki 1999, nom. inval. (art. 5).

**Nomenclatural type:** Hadač et al., 1988, Tab. 3, rel. 4, holotypus.

**Diagnostic taxa:** *Achillea stricta*, *Calamagrostis arundinacea*, *Carex umbrosa*, *Centaurea carpatica*, *Cruciata glabra*, *Dianthus compactus*, *Hieracium umbellatum*, *Lilium martagon*, *Pyrethrum clusii*, *Stellaria graminea*, *Thesium alpinum*, *Thymus pulegioides*, *Viola dacica* (Table 1).

**Constant taxa:** *Agrostis capillaris*, *Calamagrostis arundinacea*, *Cruciata glabra*, *Deschampsia cespitosa*, *Dianthus compactus*, *Festuca rubra*, *Gymnadenia conopsea*, *Hypericum maculatum*, *Luzula luzuloides*, *Potentilla erecta*, *Pyrethrum clusii*, *Scorzonera rosea*, *Stellaria graminea*, *Thymus pulegioides*, *Vaccinium myrtillus*, *Vitis-idaea*.

**Distribution in the Ukrainian Carpathians:** Beskydy, Gorgany, Hryniava Mts (Fig. 2, b).

**References:** Hadač et al., 1988, 1995; Winnicki, 1999; Kliment, Jarolimek, 2003; Kliment et al., 2004; Stoyko et al., 2007.

### *Calamagrostion villosae* Pawłowski et al. 1928

Medium-rich or species-poor chionophilous tall-grass subalpine vegetation.

**Syn.:** *Aconition firmi* Krajina 1933 (syntax. syn.), *Calamagrostion villosae* Pawłowski 1928 (art. 8), *Deschampsion caespitosae* Borza 1934 (art. 29c, 31), *Phleo alpini-Deschampsion caespitosae* (Borza 1934) Csürös et al. 1985 (art. 29c), *Poo chaixii-Deschampsion cespitosae* Jeník et al. 1980 (art. 29c), *Trisetion fusci* Krajina 1933 (syntax. syn.).

### *Poo chaixii-Deschampsietum cespitosae* Pawłowski et Walas 1949

Chionophilous tall-grass subalpine pastures (“polonya”) often dominated by *Deschampsia cespitosa*, on deep wet humus-rich loamy acidic alluvium in depressions, snowfields, near rivers and streams in mosaic with *Pinus mugo* scrub on slightly to moderately (10–20°) steep slopes at elevations between 1300 and 1900 m. a.s.l.

**Syn.:** *Aconito firmi-Deschampsietum alpicolae* (Krajina 1933) Hadač in Mucina et Maglocký 1985 (art. 2b),

*Deschampsietum caespitosae* Krajina 1933 (art. 31), *Diantho compacti-Hypericetum maculati* Winnicki 1999, nom. invalid. (art. 5), *Gentiano asclepiadeae-Acetosum carpaticae* Hadač et al. 1988 (syntax. syn.), *Phleo alpini-Deschampsietum caespitosae* (Krajina 1933) Coldea 1983 (syntax. syn.), *Rumici-Deschampsietum caespitosae* Csűrös et al. 1985 (syntax. syn.), *Scorzonero roseae-Poetum chaixii* Hadač et al. 1995, nom. invalid. (art. 5), *Trollio altissimi-Knautietum dipsacifoliae* Winnicki 1999, nom. invalid. (art. 5), *Viola dacicae-Deschampsietum caespitosae* Raťiu et al. 1983 (syntax. syn.).

**Nomenclatural type:** Pawłowski et Walas, 1949: 142 – 144, Table 10, rel. 3, lectotypus (Michl et al., 2010).

**Diagnostic taxa:** *Cardaminopsis halleri*, *Carex ovalis*, *Cerastium fontanum*, *Crocus heuffelianus*, *Deschampsia cespitosa* (dom., opt.), *Hieracium aurantiacum*, *Omalotheca norvegica*, *Phleum alpinum*, *Poa chaixii*, *Polygonum bistorta*, *Potentilla aurea*, *Ranunculus acris*, *Trifolium repens*, *Veronica officinalis*, *Viola declinata* (Table 1).

**Constant taxa:** *Achillea stricta*, *Antoxanthum odoratum*, *Campanula abietina*, *Deschampsia cespitosa*, *Festuca picta*, *Homogyne alpina*, *Hypericum alpigenum*, *H. maculatum*, *Luzula luzuloides*, *Nardus stricta*, *Phleum alpinum*, *Poa chaixii*, *Potentilla aurea*, *Vaccinium myrtillus*.

**Distribution in the Ukrainian Carpathians:** Chornohora, Svydovets', Chyvchyny, Hryniava Mts, Marmarosh Mts, Gorgany, Beskydy (Fig. 2, c).

**References:** Krajina, 1933; Pawłowski & Walas, 1949; Coldea, 1983, 1991; Csűrös et al., 1985; Hadač et al., 1988, 1995; Sanda et al., 1997; Kliment et al., 2007, 2010; Michl et al., 2010.

#### *Hyperico alpigeni-Calamagrostietum villosae* Pawłowski & Walas 1949 nom. invers. propos.

Relatively species rich communities dominated by *Calamagrostis villosa* on the intermediate and the upper parts of open steep slopes and depressions with deep and prolonged snow cover on well-drained siliceous bedrock with humus-poor skeletal soils in the subalpine belt (1500–1900 m).

**Syn.:** *Hyperico grisebachii-Calamagrostietum villosae* Pawłowski et Walas 1949 corr. Kricsfalussy et Malynovski 2000 (art. 34).

**Original name form:** Association à *Calamagrostis villosa* et *Hypericum alpigenum* = *Calamagrostidetum pocuticum* Pawł. et Wal. (Pawł 1936).

**Nomen inversum propositum:** according art. 10 b, 14, 42 ICPN.

**Nomenclatural type:** Pawłowski & Walas 1949: 140, Table 9, rel. 1, lectotypus hoc loco.

**Diagnostic taxa:** *Calamagrostis villosa* (dom., opt.), *Carex atrata*, *Doronicum austriacum*, *Hypericum alpigenum*, *Festuca fallax*, *Gentiana punctata*, *Geum montanum*, *Ligusticum mutellina*, *Phyteuma vagneri*, *Pulsatilla alba*, *Ranunculus platanifolius*, *Soldanella hungarica*, *Veratrum album* (Table 1).

**Constant taxa:** *Calamagrostis villosa*, *Deschampsia cespitosa*, *Homogyne alpina*, *Hypericum alpigenum*, *Ligusticum mutellina*, *Luzula luzuloides*, *Potentilla aurea*, *Soldanella hungarica*, *Solidago virgaurea*, *Vaccinium myrtillus*.

**Distribution in the Ukrainian Carpathians:** Chyvchyny, Hryniava Mts, Marmarosh Mts, Chornohora, and western Gorgany (Fig. 2, d).

**References:** Pawłowski & Walas, 1949; Coldea, 1991; Sanda et al., 1997; Malynovski & Kricsfalussy, 2000.

#### *Vaccinio myrtilli-Calamagrostietum villosae* Sillinger 1933

Extremely species-poor communities dominated by *Calamagrostis villosa* or *Luzula luzuloides* with relatively high cover of lichens and mosses, on cold open upper parts of slopes with very skeleton-rich shallow acid soils on flysch.

**Incl.:** *Calamagrostidetum villosae vaccinietosum myrtilli* Br.-Bl. 1930 (Kliment et al., 2007).

**Nomenclatural type:** Sillinger, 1933: 276, rel. 1, lectotypus (Kliment et al., 2004: 97).

**Diagnostic taxa:** *Calamagrostis villosa*, *Cladonia sp. div.*, *Cetraria islandica*, *Pinus mugo*, *Vaccinium vitis-idaea* (Table 1).

**Constant taxa:** *Calamagrostis villosa*, *Cladonia sp. div.*, *Cetraria islandica*, *Luzula luzuloides*, *Vaccinium myrtillus*, *V. vitis-idaea*.

**Distribution in the Ukrainian Carpathians:** Gorgany, Chornohora (Fig. 2, e).

**References:** Sillinger, 1933; Kliment et al., 2004, 2007.




Figure 2. Geographical distribution of the associations in the Ukrainian Carpathians based on phytosociological relevés: (a) *Festucetum carpaticae*, (b) *Achilleo strictae-Calamagrostietum arundinaceae*, (c) *Poo chaixii-Deschampsietum cespitosae*, (d) *Hyperico alpigeni-Calamagrostietum villosae*, (e) *Vaccinio myrtilli-Calamagrostietum villosae*.

Table 1. Synoptic table of subalpine tall-grass communities in the Ukrainian part of the Eastern Carpathians.

The species are represented by two indicators: modified fidelity measure expressed by the Phi-coefficient and frequency expressed in percentages (FIDELITY frequency).

Syntaxon number	1	2	3	4	5
Number of relevés	13	32	36	50	18
<b>D.s. Ass. Festucetum carpaticae, Al. Festucion carpaticae</b>					
<i>Festuca carpatica</i>	91.4 <sup>92</sup>	---	---	6	---
<i>Cirsium erisithales</i>	77.0 <sup>77</sup>	12	---	---	---
<i>Euphrasia picta</i>	73.3 <sup>62</sup>	---	---	2	---
<i>Aquilegia nigricans</i>	69.5 <sup>54</sup>	---	---	---	---
<i>Ranunculus oreophilus</i>	60.1 <sup>46</sup>	---	---	4	---
<i>Listera ovata</i>	59.8 <sup>54</sup>	12	---	---	---
<i>Scabiosa columbaria</i>	59.2 <sup>46</sup>	3	---	2	---
<i>Primula elatior</i>	50.8 <sup>62</sup>	19	8	14	---
<i>Galium anisophyllum</i>	48.1 <sup>31</sup>	---	3	---	---
<i>Heracleum sphondylium</i>	44.6 <sup>31</sup>	6	---	---	---
<i>Cortusa matthioli</i>	35.6 <sup>15</sup>	---	---	---	---
<i>Bupleurum longifolium</i>	30.9 <sup>15</sup>	3	---	---	---
<b>D.s. Ass. Achilleo strictae-Calamagrostietum arundinaceae</b>					
<i>Thymus pulegioides</i>	---	92.1 <sup>88</sup>	---	---	---
<i>Pyrethrum clusii</i> Fisch. ex Rchb.	---	83.3 <sup>78</sup>	---	4	---
<i>Dianthus compactus</i> Kit.	---	81.4 <sup>78</sup>	3	4	---
<i>Stellaria graminea</i>	---	72.0 <sup>69</sup>	11	---	---
<i>Centaurea carpatica</i> (Porcius) Porcius	---	61.8 <sup>47</sup>	3	---	---
<i>Viola dacica</i>	---	59.5 <sup>41</sup>	---	---	---
<i>Thesium alpinum</i>	---	56.9 <sup>38</sup>	---	---	---
<i>Hieracium umbellatum</i>	---	48.2 <sup>41</sup>	---	2	11
<i>Lilium martagon</i>	---	44.6 <sup>31</sup>	3	4	---
<i>Achillea stricta</i>	6.7 <sup>54</sup>	43.6 <sup>91</sup>	14.0 <sup>61</sup>	30	---
<i>Carex umbrosa</i>	---	39.5 <sup>19</sup>	---	---	---
<b>D.s. Ass. Phleo alpini-Deschampsietum cespitosae</b>					
<i>Phleum alpinum</i>	---	6	64.1 <sup>72</sup>	4.3 <sup>24</sup>	---
<i>Carex ovalis</i>	---	---	62.7 <sup>47</sup>	2	---
<i>Poa chiaixii</i>	---	19	61.7 <sup>81</sup>	6.5 <sup>32</sup>	---
<i>Polygonum bistorta</i>	---	12	60.8 <sup>58</sup>	4	---
<i>Ranunculus acris</i>	---	3	59.1 <sup>50</sup>	6	---
<i>Omalotheca norvegica</i>	---	---	54.4 <sup>50</sup>	4.1 <sup>16</sup>	---
<i>Cerastium fontanum</i>	---	8	53.7 <sup>53</sup>	10	---
<i>Crocus heuffelianus</i>	---	9	49.0 <sup>42</sup>	4	---
<i>Veronica officinalis</i>	---	3	47.5 <sup>31</sup>	---	---
<i>Cardaminopsis halleri</i>	---	3	31.5 <sup>19</sup>	4	---
<b>D.s. Ass. Hyperico alpigeni-Calamagrostietum villosae</b>					
<i>Gentiana punctata</i>	---	---	3	62.9 <sup>54</sup>	6
<i>Geum montanum</i>	---	---	8	49.5 <sup>38</sup>	---
<i>Carex atrata</i>	---	---	---	31.4 <sup>12</sup>	---
<i>Festuca fallax</i> Thuill.	---	---	---	31.4 <sup>12</sup>	---
<b>D.s. Ass. Vaccinio myrtilli-Calamagrostietum villosae</b>					
<i>Cladonia</i> sp. div.	---	---	3	4	77.3 <sup>72</sup>
<i>Cetraria islandica</i>	---	---	6	10	67.7 <sup>67</sup>
<i>Pinus mugo</i>	---	---	3	12	50.4 <sup>44</sup>
<b>D.s. Al. Calamagrostion arundinaceae</b>					
<i>Cruciata glabra</i>	25.4 <sup>54</sup>	61.9 <sup>88</sup>	11	---	---
<i>Calamagrostis arundinacea</i>	---	64.2 <sup>91</sup>	22	24	11
<i>Silene vulgaris</i>	---	47.3 <sup>59</sup>	6	16.1 <sup>34</sup>	6
<i>Knautia dipsacifolia</i>	7.6 <sup>23</sup>	47.3 <sup>53</sup>	8	2	---
<i>Ranunculus polyanthemos</i>	---	40.5 <sup>31</sup>	6	6	---
<i>Anemone nemorosa</i>	---	35.2 <sup>47</sup>	25.1 <sup>39</sup>	10	---

Syntaxon number	1	2	3	4	5
Number of relevés	13	32	36	50	18
<i>Betonica officinalis</i> L.	---	32.0 <sup>12</sup>	---	---	---
<i>Phyteuma spicatum</i>	---	32.0 <sup>12</sup>	---	---	---
<i>Avenula pubescens</i>	---	24.8 <sup>12</sup>	3	2	---
<b>D.s. Al. Calamagrostion villosae</b>					
<i>Hypericum alpinum</i> Kit.	---	15	---	33.2 <sup>61</sup>	49.4 <sup>76</sup>
<i>Potentilla aurea</i>	---	15	12	56.0 <sup>89</sup>	25.8 <sup>60</sup>
<i>Festuca picta</i>	---	---	3	57.2 <sup>78</sup>	30.4 <sup>54</sup>
<i>Campanula abietina</i> Griseb.	---	---	3	53.2 <sup>69</sup>	21.0 <sup>42</sup>
<i>Soldanella hungarica</i>	---	8	---	32.7 <sup>56</sup>	49.0 <sup>70</sup>
<i>Phyteuma vagneri</i>	---	---	3	20.0 <sup>31</sup>	41.1 <sup>46</sup>
<i>Luzula sylvatica</i>	---	8	---	22.8 <sup>31</sup>	39.0 <sup>42</sup>
<i>Leontodon croceus</i>	---	8	---	11.0 <sup>17</sup>	26.5 <sup>26</sup>
<i>Senecio subalpinus</i>	---	---	6	29.1 <sup>25</sup>	6.0 <sup>12</sup>
<i>Crepis conyzifolia</i>	---	---	---	14.3 <sup>8</sup>	13.3 <sup>8</sup>
<i>Avenula planiculmis</i>	---	---	---	16.6 <sup>6</sup>	2.0 <sup>2</sup>
<b>D.s. O. Calamagrostietalia villosae</b>					
<i>Luzula luzulooides</i>	---	31	23.7 <sup>91</sup>	6.9 <sup>75</sup>	12.3 <sup>80</sup>
<i>Deschampsia cespitosa</i>	---	8	21.5 <sup>69</sup>	52.8 <sup>100</sup>	12.7 <sup>60</sup>
<i>Ligusticum mutellina</i>	---	15	---	6.8 <sup>31</sup>	54.8 <sup>72</sup>
<i>Calamagrostis villosa</i>	15.0 <sup>69</sup>	---	19	39.8 <sup>94</sup>	34.7 <sup>89</sup>
<i>Laserbium alpinum</i> Waldst. & Kit.	23.6 <sup>38</sup>	6	11	7.9 <sup>26</sup>	---
<i>Centaurea mollis</i>	36.6 <sup>23</sup>	---	6	---	---
<i>Campanula serrata</i>	34.8 <sup>54</sup>	37.6 <sup>56</sup>	8	2	---
<i>Gentiana asclepiadea</i>	---	8	36.5 <sup>53</sup>	2.9 <sup>25</sup>	16
<b>D.s. Al. Adenostylium alliariae</b>					
<i>Doronicum austriacum</i>	---	---	3	3	56.7 <sup>44</sup>
<i>Adenostyles alliariae</i>	---	---	---	---	48.7 <sup>28</sup>
<i>Cirsium waldsteinii</i>	---	---	3	---	36.6 <sup>20</sup>
<i>Cicerbita alpina</i>	---	---	---	---	31.4 <sup>12</sup>
<b>D.s. Cl. Mulgedio-Aconitetea</b>					
<i>Geranium alpestre</i>	10.6 <sup>23</sup>	---	3	13.2 <sup>25</sup>	14.6 <sup>26</sup>
<i>Pedicularis hacquetii</i>	5.8 <sup>8</sup>	---	3	15.6 <sup>12</sup>	---
<i>Astrantia major</i>	51.9 <sup>69</sup>	25.9 <sup>47</sup>	3	4	---
<i>Polygonatum verticillatum</i>	13.5 <sup>15</sup>	19.7 <sup>19</sup>	---	6	---
<i>Chaerophyllum hirsutum</i>	---	8	8.3 <sup>12</sup>	10.8 <sup>14</sup>	6
<i>Myosotis caespitosa</i> Schultz	---	---	1.3 <sup>6</sup>	17.9 <sup>14</sup>	5.1 <sup>8</sup>
<i>Rumex alpinus</i>	---	---	3	39.1 <sup>22</sup>	---
<i>Leucanthemum waldsteinii</i>	---	23	26.7 <sup>91</sup>	25	2.1 <sup>26</sup>
<i>Athyrium distentifolium</i>	---	6.9 <sup>16</sup>	3	42.2 <sup>38</sup>	---
<i>Rumex alpestris</i>	---	16	25.1 <sup>44</sup>	38.8 <sup>56</sup>	---
<i>Veratrum album</i>	---	12	11	51.8 <sup>56</sup>	6
<i>Ranunculus platanifolius</i>	---	---	---	42.9 <sup>22</sup>	---
<i>Viola biflora</i>	---	8	---	38.1 <sup>28</sup>	---
<i>Stellaria nemorum</i>	---	---	1.0 <sup>6</sup>	33.8 <sup>20</sup>	---
<i>Alnus viridis</i>	---	---	3	32.2 <sup>16</sup>	---
<i>Senecio fuchsii</i> C.C.Gmel	---	---	19	15.3 <sup>19</sup>	26.1 <sup>26</sup>
<i>Pulmonaria filarszkyana</i>	---	---	3	2.1 <sup>3</sup>	20.1 <sup>8</sup>
<i>Silene dioica</i>	---	---	8.6 <sup>8</sup>	17.3 <sup>12</sup>	---
<i>Thalictrum aquilegiifolium</i>	---	3.1 <sup>3</sup>	---	19.6 <sup>8</sup>	---
<b>D.s. Al. Calamagrostion variae</b>					
<i>Epipactis atrorubens</i>	35.6 <sup>15</sup>	---	---	---	---
<i>Acinos baumgartneri</i>	25.0 <sup>8</sup>	---	---	---	---
<b>D.s. Cl. Elyno-Seslerietea</b>					
<i>Thymus pulcherimus</i>	61.2 <sup>46</sup>	---	3	---	---
<i>Festuca amethystina</i>	44.0 <sup>23</sup>	---	---	---	---
<i>Allium senescens</i>	44.0 <sup>23</sup>	---	---	---	---
<i>Gentianopsis ciliata</i> (L.) Ma	35.6 <sup>15</sup>	---	---	---	---

© 2012, Moshkov et al.

Syntaxon number	1	2	3	4	5		Syntaxon number	1	2	3	4	5			
Number of relevés	13	32	36	50	18		Number of relevés	13	32	36	50	18			
<i>Galium suberectum</i> Klokov	35.6 <sup>15</sup>	---	---	---	---		<b>Other species</b>								
<i>Linum extraaxillare</i> Kit.	35.6 <sup>15</sup>	---	---	---	---		<i>Vaccinium vitis-idaea</i>	---	46	20.1 <sup>69</sup>	---	19	---	20	40.3 <sup>89</sup>
<i>Gentiana lacinata</i>	35.6 <sup>15</sup>	---	---	---	---		<i>Vaccinium myrtillus</i>	---	15	3.6 <sup>72</sup>	7.0 <sup>75</sup>	18.8 <sup>86</sup>	27.9 <sup>94</sup>		
<i>Astragalus krajinae</i> Domin	25.0 <sup>8</sup>	---	---	---	---		<i>Solidago virgaurea</i>	---	15	5.0 <sup>47</sup>	---	33	22.4 <sup>64</sup>	8.2 <sup>50</sup>	
<i>Anemone narcissiflora</i> L.	25.0 <sup>8</sup>	---	---	---	---		<i>Polytrichum</i> sp.	---	---	44.7 <sup>72</sup>	7.6 <sup>38</sup>	14.6 <sup>44</sup>			
<i>Aster alpinus</i>	25.0 <sup>8</sup>	---	---	---	---		<i>Pleurozium schreberi</i>	---	---	42.4 <sup>50</sup>	---	16	6.0 <sup>22</sup>		
<i>Anthyllis vulneraria</i>	25.0 <sup>8</sup>	---	---	---	---		<i>Linum catharticum</i>	57.7 <sup>38</sup>	---	---	---	---	---		
<i>Phyteuma orbiculare</i>	24.6 <sup>15</sup>	---	---	2.7 <sup>6</sup>	---		<i>Rhinanthus alpinus</i>	49.3 <sup>38</sup>	---	3	---	6	---		
<i>Poa rehmannii</i>	20.9 <sup>8</sup>	---	---	---	2		<i>Parnassia palustris</i>	48.8 <sup>54</sup>	---	12	---	11	8		
<i>Carex sempervirens</i>	16.0 <sup>23</sup>	---	10.5 <sup>19</sup>	11.3 <sup>20</sup>	---		<i>Trifolium alpestre</i>	47.7 <sup>31</sup>	---	3	---	---	---		
<i>Sesleria bielzii</i>	15.2 <sup>8</sup>	---	---	10.0 <sup>6</sup>	---		<i>Euphorbia carnolica</i>	41.3 <sup>46</sup>	---	16	---	14	4		
<i>Carduus kerneri</i>	8.1 <sup>8</sup>	---	3	16.4 <sup>11</sup>	---		<i>Fragaria vesca</i>	40.4 <sup>23</sup>	---	3	---	3	---		
<b>D.s. Cl. Asplenietea</b>															
<i>Veronica urticifolia</i>	54.6 <sup>38</sup>	---	3	---	---		<i>Alchemilla monticola</i>	39.6 <sup>46</sup>	7.0 <sup>22</sup>	---	11	4	---		
<i>Valeriana tripteris</i>	38.1 <sup>38</sup>	---	---	19.6 <sup>26</sup>	---		<i>Gentianella lutescens</i>	32.5 <sup>15</sup>	---	3	---	2	---		
<i>Silene dubia</i> Herbich	30.9 <sup>15</sup>	---	3	---	---		<i>Phyteuma tetramerum</i>	30.9 <sup>15</sup>	---	3	---	3	---		
<b>D.s. Cl. Thlaspietea</b>															
<i>Saxifraga paniculata</i> Mill.	51.2 <sup>31</sup>	---	---	---	---		<i>Centaurea maramarosiensis</i>	30.9 <sup>15</sup>	---	3	---	3	---		
<i>Rumex scutatus</i>	35.6 <sup>15</sup>	---	---	---	---		<i>Salix silesiaca</i>	15.2 <sup>23</sup>	---	6	---	3	10	14.0 <sup>22</sup>	
<i>Rhodiola rosea</i>	20.2 <sup>15</sup>	---	---	17.3 <sup>14</sup>	---		<i>Picea abies</i>	5.5 <sup>31</sup>	---	16	---	25	14	21.1 <sup>44</sup>	
<b>D.s. Al. Cynosurion, Cl. Molinio-Arrhenatheretea</b>															
<i>Lotus corniculatus</i>	49.4 <sup>46</sup>	8.6 <sup>19</sup>	---	---	---		<i>Juniperus sibirica</i> Burgsd.	4.1 <sup>23</sup>	---	3	6.5 <sup>25</sup>	---	20	10.0 <sup>28</sup>	
<i>Trifolium pratense</i>	15.7 <sup>15</sup>	10.1 <sup>12</sup>	2.1 <sup>8</sup>	---	---		<i>Melampyrum saxosum</i>	48.0 <sup>54</sup>	18.2 <sup>31</sup>	---	2	---	---		
<i>Briza media</i>	---	8	43.8 <sup>31</sup>	---	---		<i>Gymnadenia conopsea</i>	35.8 <sup>62</sup>	50.6 <sup>75</sup>	---	3	6	---		
<i>Centaurea jacea</i>	---	---	42.8 <sup>22</sup>	---	---		<i>Carlina acaulis</i>	11.4 <sup>23</sup>	44.8 <sup>47</sup>	---	3	2	---		
<i>Dactylis glomerata</i>	---	---	36.0 <sup>22</sup>	6	---		<i>Vaccinium uliginosum</i>	1.9 <sup>8</sup>	11.5 <sup>12</sup>	---	6	2.5 <sup>8</sup>	---		
<i>Polygala vulgaris</i>	---	---	35.9 <sup>16</sup>	---	---		<i>Galium verum</i>	13.9 <sup>15</sup>	8.6 <sup>12</sup>	---	6	6	---		
<i>Rhinanthus minor</i>	---	---	32.0 <sup>12</sup>	---	---		<i>Campanula glomerata</i>	30.5 <sup>38</sup>	24.9 <sup>34</sup>	---	6	2	---		
<i>Festuca pratensis</i>	---	---	32.0 <sup>12</sup>	---	---		<i>Trollius europaeus</i>	24.7 <sup>31</sup>	25.4 <sup>31</sup>	---	3	4	---		
<i>Lathyrus pratensis</i>	---	---	31.4 <sup>22</sup>	3.9 <sup>8</sup>	2		<i>Filipendula ulmaria</i>	4.9 <sup>8</sup>	8.6 <sup>9</sup>	6.3 <sup>8</sup>	2	---	---		
<i>Carex pallescens</i>	---	---	30.5 <sup>28</sup>	15.9 <sup>19</sup>	2		<i>Maianthemum bifolium</i>	33.7 <sup>23</sup>	5.9 <sup>9</sup>	---	3	3	---		
<i>Prunella vulgaris</i>	---	8	28.7 <sup>25</sup>	4.2 <sup>11</sup>	---		<i>Stachys alpina</i>	20.7 <sup>15</sup>	7.8 <sup>9</sup>	---	4	4	---		
<i>Campanula patula</i>	---	---	18.6 <sup>12</sup>	3	7.9 <sup>8</sup>		<i>Mercurialis perennis</i>	19.0 <sup>8</sup>	3.3 <sup>3</sup>	---	3	3	---		
<i>Rumex acetosa</i>	---	---	16.7 <sup>6</sup>	3.7 <sup>3</sup>	---		<i>Galium schultesii</i>	16.2 <sup>8</sup>	1.8 <sup>3</sup>	---	2	2	---		
<i>Leontodon autumnalis</i>	---	---	5.3 <sup>3</sup>	14.6 <sup>6</sup>	---		<i>Ajuga reptans</i>	14.9 <sup>8</sup>	10.5 <sup>6</sup>	---	3	3	---		
<i>Alopecurus pratensis</i>	---	---	5.3 <sup>3</sup>	14.6 <sup>6</sup>	---		<i>Gentianella amarella</i>	14.9 <sup>8</sup>	10.5 <sup>6</sup>	---	3	3	---		
<i>Trifolium repens</i>	---	8	---	9	46.5 <sup>42</sup>			<i>Betula pendula</i>	16.2 <sup>8</sup>	1.8 <sup>3</sup>	---	2	2	---	
<i>Anthoxanthum odoratum</i>	---	---	25	36.7 <sup>69</sup>	22.6 <sup>56</sup>	22		<i>Aposeris foetida</i>	19.6 <sup>31</sup>	---	12	26.9 <sup>36</sup>	2	---	
<i>Poa pratensis</i>	---	---	---	30.2 <sup>11</sup>	---			<i>Antennaria dioica</i>	10.7 <sup>8</sup>	---	3	5.0 <sup>6</sup>	2	---	
<i>Vicia sepium</i>	---	---	3	26.4 <sup>14</sup>	2			<i>Paris quadrifolia</i>	17.7 <sup>8</sup>	---	3	5.5 <sup>4</sup>	4	---	
<b>D.s. Al. Nardo-Agrostion, Cl. Calluno-Ulicetea</b>															
<i>Hypochoeris uniflora</i>	---	---	50.3 <sup>66</sup>	17	5.7 <sup>28</sup>	6		<i>Euphrasia tatrae</i> Wettst.	15.2 <sup>8</sup>	---	3	10.0 <sup>6</sup>	6	---	
<i>Potentilla erecta</i>	---	8	56.2 <sup>75</sup>	14.9 <sup>39</sup>	2	6		<i>Heracleum carpaticum</i>	13.8 <sup>15</sup>	---	8	14.9 <sup>16</sup>	16	---	
<i>Scorzonera rosea</i> Waldst. & Kit.	---	---	47.6 <sup>72</sup>	17.3 <sup>44</sup>	22	6		<i>Festuca versicolor</i>	26.6 <sup>15</sup>	---	3	4	4	---	
<i>Agrostis capillaris</i>	---	---	42.8 <sup>56</sup>	35.1 <sup>50</sup>	---	---		<i>Rosa pendulina</i>	32.5 <sup>15</sup>	---	3	2	2	---	
<i>Nardus stricta</i>	---	---	25.3 <sup>41</sup>	40.4 <sup>53</sup>	8	---		<i>Galium mollugo</i>	20.9 <sup>8</sup>	---	3	2	2	---	
<i>Viola declinata</i>	---	---	11.9 <sup>25</sup>	45.9 <sup>50</sup>	6	---		<i>Carex ornithopoda</i>	20.9 <sup>8</sup>	---	2	2	2	---	
<i>Hieracium aurantiacum</i>	---	---	1.0 <sup>16</sup>	53.2 <sup>53</sup>	6	---		<i>Luzula sudetica</i>	44.0 <sup>23</sup>	---	3	3	3	---	
<i>Luzula multiflora</i>	---	---	4.7 <sup>12</sup>	35.2 <sup>31</sup>	6	---		<i>Thymus alpestris</i>	18.5 <sup>38</sup>	6	32.2 <sup>50</sup>	20	---		
<i>Arnica montana</i>	---	15	7.2 <sup>28</sup>	10.1 <sup>31</sup>	20	17			<i>Brachythecium</i> sp.	8.5 <sup>8</sup>	---	3.2 <sup>6</sup>	9.3 <sup>8</sup>	---	
<i>Carex pilulifera</i>	---	---	9.0 <sup>3</sup>	7.4 <sup>3</sup>	---	---			<i>Hypericum maculatum</i>	8	45.3 <sup>78</sup>	33.3 <sup>67</sup>	22	---	
<b>D.s. Cl. Caricetea curvulae</b>															
<i>Avenula versicolor</i>	10.6 <sup>8</sup>	---	3	11.4 <sup>8</sup>	---				<i>Festuca rubra</i>	8	43.4 <sup>62</sup>	22.6 <sup>44</sup>	10	---	
<i>Festuca airoides</i>	7.0 <sup>8</sup>	---	21.6 <sup>14</sup>	2	---				<i>Veronica chamaedrys</i>	---	30.5 <sup>31</sup>	20.8 <sup>25</sup>	2	---	
<i>Homogyne alpina</i>	---	15	50	25.7 <sup>78</sup>	31.9 <sup>84</sup>	33					5.7 <sup>6</sup>	25.1 <sup>14</sup>	4	---	
<i>Juncus trifidus</i>	---	---	---	42.9 <sup>22</sup>	---						2.8 <sup>3</sup>	20.2 <sup>8</sup>	2	---	
<i>Pulsatilla alba</i>	---	---	---	40.8 <sup>20</sup>	---						2.8 <sup>3</sup>	20.2 <sup>8</sup>	3	---	
<i>Hieracium alpinum</i>	---	---	6	36.8 <sup>38</sup>	13.4 <sup>22</sup>	↑					9.0 <sup>3</sup>	7.4 <sup>3</sup>	3	---	
<i>Huperzia selago</i>	---	---	---	14.4 <sup>10</sup>	17.1 <sup>11</sup>	↓					9.0 <sup>3</sup>	7.4 <sup>3</sup>	4	---	

Syntaxon number Number of relevés	1 13	2 32	3 36	4 50	5 18
<i>Ranunculus repens</i>	---	---	22.0 <sup>8</sup>	---	---
<i>Campanula cervicaria</i>	---	---	16.6 <sup>6</sup>	2.0 <sup>2</sup>	---
<i>Hylocomium</i> sp.	---	---	14.3 <sup>19</sup>	24.9 <sup>26</sup>	---
<i>Oxalis acetosella</i>	---	---	19.2 <sup>17</sup>	21.8 <sup>18</sup>	---
<i>Urtica dioica</i>	---	---	16.6 <sup>6</sup>	2.0 <sup>2</sup>	---
<i>Rhytidadelphus squarrosum</i>	---	---	39.0 <sup>42</sup>	22.4 <sup>30</sup>	---
<i>Leontodon hispidus</i>	---	6	37.8 <sup>33</sup>	2.8 <sup>12</sup>	---
<i>Polygonatum alpinum</i>	---	---	10.8 <sup>6</sup>	12.3 <sup>6</sup>	---
<i>Phragmites connectilis</i>	---	---	6.1 <sup>3</sup>	11.4 <sup>4</sup>	---
<i>Mnium</i> sp.	---	---	4.2 <sup>6</sup>	26.0 <sup>14</sup>	---
<i>Caltha laeta</i> Schott, Nyman & Kotschy	---	---	9.4 <sup>3</sup>	5.4 <sup>2</sup>	---
<i>Cardamine pratensis</i>	---	---	9.4 <sup>3</sup>	5.4 <sup>2</sup>	---
<i>Hieracium prenanthoides</i>	---	---	9.4 <sup>3</sup>	5.4 <sup>2</sup>	---
<i>Rubus idaeus</i>	---	---	5.6 <sup>6</sup>	23.1 <sup>12</sup>	---
<i>Coeloglossum viride</i>	---	---	3.9 <sup>3</sup>	16.2 <sup>6</sup>	---
<i>Crepis paludosa</i>	---	---	9.4 <sup>3</sup>	5.4 <sup>2</sup>	---
<i>Carex sylvatica</i>	---	---	9.4 <sup>3</sup>	5.4 <sup>2</sup>	---
<i>Epilobium angustifolium</i>	---	3	---	28.2 <sup>18</sup>	---
<i>Rhododendron kotschyi</i>	---	---	---	44.9 <sup>24</sup>	---
<i>Campanula rotundifolia</i>	8	---	6	33.7 <sup>26</sup>	---
<i>Myosotis alpestris</i>	---	---	3	32.2 <sup>16</sup>	---
<i>Rhytidadelphus triquetrus</i>	---	---	6	36.1 <sup>22</sup>	---
<i>Sphagnum</i> sp.	---	---	4.5 <sup>6</sup>	10.9 <sup>8</sup>	4.5 <sup>6</sup>
<i>Melampyrum herbichii</i>	---	9	---	3.8 <sup>16</sup>	29.2 <sup>33</sup>
<i>Sorbus aucuparia</i>	---	---	---	3.5 <sup>6</sup>	29.2 <sup>17</sup>
<i>Dicranum</i> sp.	---	---	6	---	30.8 <sup>22</sup>

#### Taxa with low frequency occurring in one association only:

1: *Abies alba*, *Spiraea chamaedryfolia* (35.6<sup>15</sup>); *Asplenium viride*, *Agrostis rupestris*, *Cerastium lanatum* Lam., *Daphne mezereum*, *Epipactis helleborine*, *Hieracium villosum* (25.0<sup>8</sup>).

2: *Allium victorialis*, *Phleum pratense*, *Plantago lanceolata*, *Thalictrum minus* (27.6<sup>9</sup>); *Botrychium lunaria*, *Brachypodium pinnatum*, *Centaurea kotschyana*, *Dianthus deltoides*, *Digitalis grandiflora*, *Euphrasia rostkoviana*, *Gymnadenia odoratissima*, *Knautia arvensis*, *Trifolium medium*, *T. montanum* (22.5<sup>6</sup>); *Aconitum lasiocarpum* Gayer, *A. moldanicum* Hacq., *Aegopodium podagraria*, *Athyrium filix-femina*, *Campanula persicifolia*, *Carex brizoides*, *C. hirta*, *Chamaespartium saggittale*, *Cirsium oleraceum*, *Cynosurus cristatus*, *Dactylorhiza sambucina*, *Dianthus carthusianorum*, *Empetrum hermaphroditum* Hagerup, *Euphrasia stricta*, *Festuca apennina* De Not., *Genista tinctoria*, *Holcus mollis*, *Lathyrus sylvestris*, *Lysimachia vulgaris*, *Orchis mascula* subsp. *signifera*, *Origanum vulgare*, *Picris hieracioides*, *Pimpinella major*, *Polygala comosa*, *Scrophularia scopolii*, *Danthonia decumbens*, *Soldanella montana*, *Tragopogon pratensis*, *Vicia cracca*, *Viola riviniana* (15.9<sup>3</sup>).

3: *Rumex acetosella* (33.8<sup>14</sup>); *Euphorbia amygdaloides* (26.0<sup>8</sup>); *Hieracium pilosella*, *Pseudorchis albida*, *Luzula campestris*, *Poa alpina*, *Taraxacum officinale*, *Veronica serpyllifolia* (21.2<sup>6</sup>); *Achillea cartilaginea*, *Cardamine flexuosa*, *Carex nigra*, *Diphasiastrum alpinum*, *Juncus effusus*, *Lychnis flos-cuculi*, *Lycopodium clavatum*, *Pedicularis verticillata*, *Silene pusilla*, *Sympyton cordatum* (14.9<sup>3</sup>).

4: *Hieracium* sp. (33.9<sup>14</sup>); *Aconitum nanum* (Baumg.) Simonk., *Luzula alpinopilosa*, *Milium effusum*, *Ranunculus carpaticus* (28.6<sup>10</sup>); *Campanula alpina*, *Empetrum nigrum*, *Hieracium atratum*, *Poa deylii* Chrtk & V.Jirásek, *Rumex obtusifolius*, *Valeriana sambucifolia* J.C.Mikan (22.0<sup>6</sup>); *Dryopteris dilatata*, *Pedicularis sylvatica*, *Veronica baumgartneri* (25.5<sup>8</sup>); *Aconitum variegatum*, *Bartsia alpina*, *Chrysosplenium alpinum*, *Eurychium pilferum*, *Lamiastrum galeobdolon*, *Pohlia nutans* (18.0<sup>4</sup>); *Achillea lingulata*, *Adoxa*

*moschatellina*, *Allium ursinum*, *Angelica archangelica*, *Atrichum* sp., *Bryum* sp., *Chrysosplenium alternifolium*, *Clematis alpina*, *Cystopteris sudetica*, *Dentaria glandulosa* Waldst. & Kit., *Dryopteris filix-mas*, *D. carthusiana*, *Epilobium montanum*, *Euphrasia salisburgensis*, *Gentiana acaulis*, *Hieracium grandidens*, *H. nigrescens*, *Lloydia serotina*, *Potentilla reptans*, *Ribes alpinum*, *Rhinanthus angustifolius* (12.7<sup>2</sup>).

5: *Pinus cembra* (21.2<sup>6</sup>).

#### Syntaxon numbers:

1 – *Festucetum carpaticae*, 2 – *Achilleo strictae-Calamagrostietum arundinaceae*, 3 – *Poo chaixii-Deschampsietum cespitosae*, 4 – *Hyperico alpigeni-Calamagrostietum villosae*, 5 – *Vaccinio myrtilli-Calamagrostietum villosae*.

#### Sources of the relevés:

1: Malynovski & Kricsfalussy, 2000, P. 93, Table 20, rel. 8-12 (5 rel.), Chornei I. & Tokaryuk A., 2009, unpubl. (8 rel.);

2: Hadač et al., 1995, P. 687 (1 rel.), Kvakovska I., 2009, unpubl. (21 rel.), Tokaryuk A., 2009, unpubl. (5 rel.), Burlaka M., 2008., unpubl. (5 rel.);

3: Pawłowski & Walas, 1949, Table X, rel. 1-16 (16 rel.), Malynovski & Kricsfalussy, 2000, P. 160, Table 44, rel. 1-11 (11 rel.), Malynovski & Kricsfalussy, 2000, P. 163, Table 45, rel. 1-8 (8 rel.), Solomakha et al., 2004, P. 79, Table 3.15, rel. 4 (1 rel.);

4: Pawłowski & Walas, 1949, Table IX, rel. 1-20 (20 rel.), Malynovski & Kricsfalussy, 2000, P. 156, Table 43, rel. 1-15 (15 rel.), Burlaka M., 2008, unpubl. (10 rel.), Iakushenko D., 2008, unpubl. (5 rel.);

5: Burlaka M., 2008, unpubl. (13 rel.), Iakushenko D., 2007, unpubl. (5 rel.).

## DISCUSSION

The results show clear differences between the five groups of subalpine tall-grass communities (Table 1). However, their positions within the order *Calamagrostietalia villosae* remain still rather uncertain.

The syntaxonomical distinctiveness of the alliance *Festucion carpaticae* is questioned by some authors. In the most recent survey, Michl et al. (2010) merged this alliance with *Calamagrostion villosae*. An intermediate position of this syntaxon between subalpine tall-grass vegetation dominated by *Calamagrostis arundinacea* or *C. villosa* and calciphilous communities belonging to the class *Elyno-Sesslerietea* Br.-Bl. 1948 was repeatedly mentioned (Kliment et al., 2004; Kliment & Valachovič, 2007). In the data analyzed in our work, relevés of the association *Festucetum carpaticae* contain a lot of species of the class *Elyno-Sesslerietea*; some

of them attain high fidelity values (*Thymus pulcherrimus*, *Festuca amethystina*, *Allium montanum*, *Gallium suberectum*, *Gentiana laciñata*, *Linum extraaxillare* etc) (Table 1). Moreover, some relevés classified as representatives of *Thymo-Festucetum amethystinae festucetosum carpaticae* Kricsfalussy et Malynovski 2000 within the class *Elyno-Seslerietea* (Malinovski & Kricsfalussy, 2000: 93-101), have been included by us into *Festucetum carpaticae*. We consider the alliance *Festucion carpaticae* as a separate unit because of its significant floristic and ecological peculiarity within the order *Calamagrostietalia villosae*.

The association *Festucetum carpaticae* is reported for Ukraine for the first time by Domin (1930) from Svydovets'. The corresponding relevés were cited by Bělohlávková, Fišerová (1989: 4, Table 1, rel. 11, only data of the taxa presence without semi quantitative values) as "not typical *Festucetum carpaticae*" (Ibid.: 22). The association was not mentioned in later surveys (Malynovski & Kricsfalussy, 2000; Solomakha, 2008); only Malynovski (2003) mentions this syntaxon without references or relevés.

Quite interesting are the syntaxonomical positions of species-rich montane to subalpine communities dominated by *Calamagrostis arundinacea*, reported for Beskydy, but known also from Chyvchyny, Marmarosh, Gorgany and Chornohora Mts (Malynovski, 1980). Available relevés belongs to only one association *Achilleo strictae-Calamagrostietum arundinaceae* Hadač et al. 1988, described from the southern slopes of the westernmost part of the Eastern Carpathians (Hadač et al., 1988). The association *Tanaceto-Calamagrostietum arundinaceae* Winnicki 1999 invalidly described from the northern slopes of the same mountain ridge (Winnicki, 1999) does not show any significant differences from the previous one and seems to be just a synonym. Considering the relatively large species pool of the Eastern Carpathians (Czopik, 1976; Tasenkevich, 2006; Chornei, 2009), the local heterogeneity of bedrocks and the peculiarities of the land-use history, we expect a higher diversity within the alliance *Calamagrostion arundinaceae* at the association or subassociation level, especially in the Marmarosh and Chyvchyny regions (cf. Kliment et al., 2007). More detailed studies both in the Romanian and the Ukrainian parts of the region should help to solve this problem within the alliance.

Regarding *Achillea stricta*, although this species exhibits a high fidelity in communities of the association *Achilleo strictae-Calamagrostietum arundinaceae*, it probably should also be treated as a regionally diagnostic species for the order *Calamagrostietalia villosae* in the Eastern Carpathians, because of its rather high constancy in other communities. Moreover *Dianthus compactus*, *Centaurea carpatica*, and *Viola dacica* clearly reflect the East Carpathian nature of this association.

The secondary origin of these communities, formed under

the influence of extensive grazing (Malynovski, 1980, 2003) is clearly seen in the considerable proportion of mesophilous meadow species associated with the "polonya" pastures of the alliance *Cynosurion cristati* Tx. 194 (*Briza media*, *Centaurea jacea*, *Dactylis glomerata*, *Polygala vulgaris*, etc.), suballiance *Polygalo-Cynosurenion* Jurko 1974 (*Carlina acaulis*, *Campamula glomerata*, *Gymnadenia conopsea*, etc.), and the alliance *Nardo-Agrostion tenuis* Sillinger 1933 (*Hypochoeris uniflora*, *Scorzonera rosea*, *Agrostis capillaris*, etc.) (Table 1).

The association *Bupleuro-Calamagrostietum arundinacea* (Zlatník 1928) Jeník 1961, is mentioned by Malynovski (2003) without any references or relevés. The occurrence of this associations in the Ukrainian Carpathians seems to be quite plausible in "polonyas" of the montane belt in Transcarpathia, but this suggestion still needs confirmation. The other communities with a significant cover of *Calamagrostis arundinacea* are mentioned for the montane belt in Gorgany and Chornohora mountains. They are regarded as belonging to the association *Calamagrostio-Spireetum chamaedryfoliae* Resmeritā et Csürös 1966, ranked into the alliance *Calamagrostion arundinaceae*. These communities develop on siliceous flysch on steep (30-45°) slopes, rock shelves and cracks with shallow skeletal soils (Klimuk et al., 2006; Iakushenko, 2007). Considering the dominance of shrubs species (*Lonicera xylosteum*, *Rosa pendulina*, *Spiraea chamaedryfolia*, etc.) we removed such relevés from the present analysis.

Mesic and wet subalpine tall-grass pastures dominated by *Deschampsia cespitosa* are very common in the Ukrainian Carpathians due to earlier disturbance. Although their distribution, dynamic and ecological peculiarities were previously studied (Iermachenko, 1962; Malynovski, 1980), their syntaxonomical position is still ambiguous and doubtful.

Two associations, namely *Poo chaixii-Deschampsietum cespitosae* and *Phleo alpini-Deschampsietum caespitosae*, are reported for the subalpine belt of the Ukrainian Carpathians (Malynovski & Kricsfalussy, 2000; Solomakha, 2008). They belong either to the alliance *Calamagrostion villosae* (Michl et al., 2010), or to the alliance *Trisetion fuscii* (Kliment et al., 2010). In the analyzed dataset we found only few diagnostic species for the alliance *Trisetion fuscii*. All of them, except *Cerastium fontanum*, exhibit insignificant fidelity values. Although the species *Festuca picta*, *Hypericum alpinum*, *Campanula abietina*, *Soldanella hungarica*, *Phyteuma vagneri* seem to be diagnostic for the alliance *Calamagrostion villosae*, they are common both in *Phleo alpini-Deschampsietum* and in communities that undoubtedly belong to the *Calamagrostion villosae* (Table 1). So, following Michl et al. (2010), we consider the alliance *Trisetion fuscii* as syntaxonomical synonym of the alliance *Calamagrostion villosae*.

Though the authors analyzed mainly relevés from the Sudeten Mts (Michl et al., 2010: 153), they used the name *Poo chaixii-Deschampsietum cespitosae* – the association originally described from the Eastern Carpathians within the alliance *Calamagrostion villosae* (Pawłowski & Walas, 1949). We did not find any significant differences in floristic composition between the phytosociological relevés of both associations collected in the different parts of our study area (from Czywczyny to Beskydy). Thus, we suggest that *Phleo alpini-Deschampsietum caespitosae* is a syntaxonomical synonym of the *Poo chaixii-Deschampsietum cespitosae*. In this respect we should reject the name *Phleo alpini-Deschampsietum caespitosae*.

Communities dominated by *Calamagrostis villosa* belong to two associations. As mentioned above, the original name of the association *Hyperico alpigeni-Calamagrostietum villosae* was “Association à *Calamagrostis villosa* et *Hypericum alpinum* = *Calamagrostidetum pocuticum* Pawł. et Wal. (Pawł 1936)” (Pawłowski & Walas, 1949). As the second name should be rejected according to art. 34 ICPN, the first name should be corrected according to art. 10 b, 14 ICPN.

Currently, *Hypericum alpinum* Kit. is considered to be the synonym of an accepted name *Hypericum richeri* Vill. subsp. *grisebachii* (Boiss.) Nyman (Tutin et al., 1968; Ciocârlan, 2009). However, since the name *Hypericum alpinum* Kit. was validly published in 1863 (Sennikov, 1996), we believe that the association name was unreasonably (art. 34 ICPN) changed by Malynovski & Kricsfalussy (2000) on *Hyperico grisebachii-Calamagrostietum villosae* Pawł. et Wal. 1949 corr. Kricsfalussy et Malynovski 2000. *Hypericum grisebachii* Boiss is also considered to be the synonym of *Hypericum richeri* Vill. subsp. *grisebachii* (Boiss.) Nyman (Tutin et al., 1968). Consequently, the form *Hyperico grisebachii-Calamagrostietum villosae* Pawł. et Wal. 1949 used by Romanian researchers (Coldea, 1991; Sanda et al., 1997) should be also changed.

Thus, according to art. 42 we propose the following correct form of the name: *Hyperico alpigeni-Calamagrostietum villosae* Pawłowski et Walas 1949 nom. invers. propos.

The last association, *Vaccinio myrtilli-Calamagrostietum villosae*, distinguished for the Ukrainian part of the Eastern Carpathians for the first time (cf. Malynovski, Kricsfalussy, 2002). These species-poor communities with abundant cover of lichens and mosses dominated by *Calamagrostis villosa* or *Luzula luzuloides* are definitively transitional between tall-grass *Calamagrostietalia villosae* communities, on one side, and *Loiseleurio-Vaccinietea* Eggler ex Schubert 1960 and *Caricetea curvulae* Br.-Bl. 1948 communities, on the other (Table 1). Stands of this association occur mainly in the Gorgany Mts. More investigations are needed to clarify their syntaxonomical positions.

## CONCLUSIONS

Subalpine tall-grass communities in the Ukrainian part of the Eastern Carpathians are represented by five associations: *Festucetum carpaticae* (species-rich chionophilous tall-grass communities on moist carbonate soils), *Achilleo strictae-Calamagrostietum arundinaceae* (species-rich moderate thermophilous tall-grass communities in relatively warm and dry habitats on flysch), *Poo chaixii-Deschampsietum cespitosae* (chionophilous tall-grass subalpine pastures dominated by *Deschampsia cespitosa* on deep wet humus-rich acidic soils), *Hyperico alpigeni-Calamagrostietum villosae* (relatively species rich communities dominated by *Calamagrostis villosa* on siliceous bedrock), and *Vaccinio myrtilli-Calamagrostietum villosae* (extremely species-poor communities on cold slopes with shallow acidic soils). The syntaxonomy of these communities can be assessed as follows:

### **MULGEDIO-ACONITETEA HADAČ ET KLIKA IN KLIKA ET HADAČ 1944**

*Calamagrostietalia villosae* Pawłowski et al. 1928.

*Festucion carpaticae* Bělohlávková et Fišerová 1989.

*Festucetum carpaticae* Domin 1925.

*Calamagrostion arundinaceae* (Luquet 1926) Oberd. 1957.

*Achilleo strictae-Calamagrostietum arundinaceae* Hadač et al. 1988.

*Calamagrostion villosae* Pawłowski et al. 1928.

*Poo chaixii-Deschampsietum cespitosae* Pawłowski et Walas 1949.

*Hyperico alpigeni-Calamagrostietum villosae* Pawłowski et Walas 1949 nom. invers. propos.

*Vaccinio myrtilli-Calamagrostietum villosae* Sillinger 1933.

## REFERENCES

- Bělohlávková R., Fišerová D., 1989. *Festucion carpaticae* alliance nova – a new alliance of tall grasslands in the high Carpathian Mountains. *Folia Geobotanica et Phytotaxonomica* 24/1, 1–24.
- Chornei I.I., 2009. Flora Chyvchyno-Hryniavskykh hir (Ukrain's'ki Karpaty) [Flora of Chyvchyno-Hryniavsky Mts]. D. Hab. Thesis, M.G. Kholodny Institute of Botany, Kyiv.

- Chytrý M., Tichý M., Holt J., Botta-Dukát Z., 2002. Determination of diagnostic species with statistical fidelity measures. *Journal of Vegetation Science* 13, 79–90.
- Ciocârlan V., 2000. Flora ilustrată a României (Pteridophyta et Spermatophyta). Editura Ceres, Bucureşti.
- Coldea Gh., 1983. Precizări sintaxonomice asupra cenozelor cu *Deschampsia caespitosa*. Studii si Cercetari de Biologie. Ser. Biologie Vegetala 35, 96–100.
- Coldea Gh. 1991. Prodrome des associations végétales des Carpates du Sud-Est (Carpates Roumaines). Documents Phytosociologiques 13, 317–359.
- Csűrös Š., Gergely I., Csűrös-Káptalan M., 1985. Studii fitocenologice asupra pajiștilor de *Deschampsia caespitosa* (L.) P. Beauv. din R. S. România. Contribuții Botanice, 141–159.
- Czopik V.I., 1976. Vysokohirna flora Ukrains'kykh Karpat [High-altitude flora of Ukrainian Carpathians]. Naukova dumka, Kyiv.
- Deyl M., 1940. Plant, soil and climate of Pop Ivan. Synecological study from Carpathian Ukraina. Opera Botanica Čechica 2, 1–290.
- Domin K., 1925. *Festucetum carpaticae* v Bielských Tatrách. Rozpr. II. Třídy České Akad. Věd. Umění 34/19, 1–25.
- Domin K., 1930. Šimanův kotel na Svidovci v Podkarpacké Rusi. Věstn. Král. Čes. Společ. Nauk, Praha, 4 (p. separ.), 1–20.
- Hadač E., Andresová J., Klescht V., 1988. Vegetace polonin v Bukovských vrších na sv. Slovensku. Preslia 60, 321–338.
- Hadač E., Stoyko S.M., Terray J., Tasenkevich L., Bural M., 1995. Notes on plant communities of the protected complex Stužhytzia – a part of the trilateral Polish-Slovakian-Ukrainian biosphere reserve “The Eastern Carpathians”. Ukrainian Botanical Journal 52/5, 686–696.
- Hennekens S.M., Schaminée J.H.J., 2001. TURBOVEG, a comprehensive data base management system for vegetation data. *Journal of Vegetation Science* 12: 589 – 591.
- Hill M.O., 1979. TWINSPLAN: A FORTRAN program for arranging multivariate data in an ordered two-way table by classification of the individuals and attributes. Section of Ecology and Systematics. Cornell University, Ithaca, NY.
- Iakushenko D., 2007. Dopovnennia do klasyfikacij vysokotrvnoji roslynnosti Ukrains'kyh Karpat [Addition to classification of tall-herbs vegetation of Ukrainian Carpathians]. Ukrainian Botanical Journal 64/3, 426–437.
- Iermachenko H.A., 1962. Geobotanicheskaja kharakteristika lugov shchuchki dernistoj v Chernogore [Geobotanical characteristic of *Deschampsia caespitosa* meadows in Chornohora]. Ph.D. Thesis, Lviv.
- Klement J., Jarolímek I., 2003. Syntaxonomical revision of the plant communities dominated by *Calamagrostis arundinacea* (alliance *Calamagrostion arundinaceae*) in Slovakia. Thaiszia – Journal of Botany 13: 135–158.
- Klement J., Jarolímek I., Kochjarová J., 2004. Spoločenstvá zväzu *Calamagrostion arundinaceae* v Bukovských vrchoch. Bull. Slov. Bot. Spoločn. 26, 137–144.
- Klement J., Jarolímek I., Šibík J., 2007. *Mulgedio-Aconitetea* Hadač et Klika in Klika 1948. In: J. Klement & M. Valachovič (Eds). Rastlinné spoločenstvá Slovenska. 4. Vysokohorská vegetácia. pp. 21–129. Veda, Bratislava.
- Klement J., Jarolímek I., Šibík J., Valachovič M., 2004. Syntaxonomy and nomenclature of the communities of the orders *Calamagrostietalia villosae* and *Adenostyletalia* in Slovakia. Thaiszia – Journal of Botany 14 (2), 93–157.
- Klement J., Šibík J., Šibíková I., Jarolímek I., Dúbravcová Z., Uhlířová J., 2010. High-altitude vegetation of the Western Carpathians – a syntaxonomical review. Biologia 65/6, 96 –989.
- Klement J., Valachovič M. (Eds), 2007. Rastlinné spoločenstvá Slovenska. 4. Vysokohorská vegetácia, pp. 388. Veda, Bratislava.
- Klimuk Iu.V., Miskevich U.D., Iakushenko D.M., Chornei I.I., Budzhak V.V., Nyporko S.O., Shpilchak M.B., Cherniavskyi M.V., Tokaryuk A.I., Oleksiv T.M., Tymchuk Ia.Ia., Solomakha V.A., Solomakha T.D., Maior R.V. 2006. Pryrodnji zapovidnyk Gorgany. Roslynnyi svit [Natural reserve Gorgany. Plant world], Fitodosiocentr, Kyiv.
- Kočí M., 2001. Subalpine tall-forb vegetation (*Mulgedio-Aconitetea*) in the Czech Republic – syntaxonomical revision. Preslia 73, 289–331.
- Kočí M., 2007. Subalpínská vysokobylinná a křovinná vegetace (*Mulgedio-Aconitetea*). In: M. Chytrý (ed.), Vegetace České republiky 1. Travinná a keříčková vegetace, pp. 91–131. Academia, Praha.
- Krajina V., 1933. Die Pflanzengesellschaften des Mlynica-Tales in den Vysoké Tatry (Hohe Tatra). Beih. Bot. Centralbl., 50 B, 1–224.
- Malynovski K.A., 1980. Vysokohirna roslynnist' Ukrains'kykh Karpat [High-altitude vegetation of Ukrainian Carpathians]. Naukova dumka, Kyiv.
- Malynovski K., 2003. Karpats'ky polonyny i polonyns'ke hospodarstvo [Carpathian poloninas and poloninians economy]. Pracy naukovogo tovarystva im. Shevchenka. Ekolohichnyi zbirnyk 12, 293–09.

- Malynovski K.A., Kricsfalusy V.V., 2000. Roslynnist' Ukrayiny. 1. Vysokohirska roslynnist' [Vegetation of the Ukraine. 1. High mountain vegetation], pp. 232. Phytosociocentre, Kyiv.
- Malynovski K.A., Kricsfalusy V.V., 2002. Roslynni uhrupovannia vysokohiria Ukrains'kykh Karpat [Plant communities of the Ukrainian Carpathian highlands]. pp. 244. Uzhorod.
- Malynovski K.A., Mirkin B.M., Ishbirdin A.R., Kricsfalusy V.V., Komendar V.I., 1992. Syntaksonomiia pryberezhno-vodnykh, bolotnykh, luchnykh, chaharnykovykh i chaharnychkovykh uhrupovan' vykohiria Ukrains'kykh Karpat [Syntaxonomy of riparian, bog, meadow, shrub and dwarf-shrub plant communities in the highlands of Ukrainian Carpathians]. Ukrainian Botanical Journal 49/4, 5–13.
- Malynovski K.A., Tsaryk I.V., 1995. Novi dla Ukrayiny syntaksony z Karpat [New for Ukraine syntaxa from the Carpathians]. Ukrainian Botanical Journal 52/5, 621–639.
- Matuszkiewicz W., 2006. Przewodnik do oznaczania zbiorowisk roślinnych Polski, Wydawnictwo Naukowe PWN, Warszawa.
- Michl T., Dengler J., Huck S., 2010. Montane-subalpine tall-herb vegetation (*Mulgedio-Aconitetea*) in central Europe: large-scale synthesis and comparison with northern Europe. Phytocoenologia 40 (2-3), 117–154.
- Pawłowski B., Walas J., 1949. Les associations des plantes vasculaires des Monts de Czywczyn. Bulletin International de l'Académie Polonaise des Sciences et des Lettres. Série B: Sciences Naturelles 1, 1–181.
- Sanda V., Öllerer K., Burescu P., 2008. Fitocenozele din Romania. Ars Docendi, București.
- Sanda V., Popescu A., Barabaș N., 1997. Cenotaxonomia și caracterizarea grupărilor vegetale din România. Biologie vegetală 14: 1–366.
- Sennikov A.I., 1996. *Hypericaceae*. In: N.M. Tzvelev (Ed) Flora Europae Orientalis. IX Vol., pp. 173–177. Sankt-Peterburg.
- Šibíková I., Šibík J., Jarolímek I., 2008. The tall-herb and tall-grass plant communities of the class *Mulgedio-Aconitetea* in the subalpine belt of the Krivánska Malá Fatra Mts (Slovakia). Hacquetia 7/2, 141–159.
- Sillinger P., 1933. Monograficka studie o vegetaci Nizkych Tater. Orbis, Praha.
- Solomakha V., Iakushenko D., Kramarets V., Milkina L., Vorontsov D., Vorobyov Ie., Voytiuk B., Vinichenko T., Kokhanets M., Solomakha I., 2004. Nacionalyi pryrodnyi park Skoliv's'ki Beskydy. Roslynnyj svit [National nature park Skoliv's'ki Beskydy. Plant world]. Fitosociocentr, Kyiv.
- Solomakha V.A., 2008. Syntaksonomiia roslynnosti Ukrayiny. Tretie nablyzhennia [Syntaxonomy of the vegetation of Ukraine. Third approximation]. Fitosociocentr, Kyiv.
- Stoyko S.M., Hadač E., Tasenkevich L., Bural M., Terray J., 2007. Uzhans'kyi natsionalnyi pryrodnyi park. Polifunktionalne znachennia [Uzhans'kyi National Nature Park. Multifunctional importance]. Merkator, Lviv.
- Swiderski W., Szafran B., 1931. Typy florystyczne połonin w Karpatach Wschodnich. Państwowy Instytut gospodarstwa wiejskiego 12, 62–114.
- Tasenkevich L.O., 2006. Prypodna flora sudynnykh roslyn Karpat, ii osoblyvosti ta genezys [Natural flora of the vascular plants of Carpathians, its peculiarities and genesis]. D. Hab. Thesis, M.G. Kholodny Institute of Botany, Kyiv.
- Tichý L., 2002. JUICE, software for vegetation classification. Journal of Vegetation Science 13, 451–453.
- Tutin T.G. et al., 1964–80. Flora Europaea, vol 1 (1964); vol 2 (1968); vol 3 (1972); vol 4 (1976); vol 5 (1980). Cambridge University Press, Cambridge.
- Unar J., Unarová M., Šmarda J., 1985. Vegetační poměry Tomanovy doliny a Žlebu spod Diery v Západních Tatrách. 2. Charakteristika přírodních poměrů a rostlinných společenstev. Folia Fac. Sci. Nat. Univ. Purkyniana Brunnense, Ser. Biol. 26/14: 5–78.
- Ustymenko P.M., Tasenkevich L.O., 2009. Uhrupovannia formacji kostrytsi karpats'koji (*Festuceta carpatica*) [Communities of the formation *Festuceta carpatica*]. In: Ya.P. Didukh (Ed) Zelena knyha Ukrayiny, pp. 258–259. Alterpress, Kyiv.
- Weber H. E., Moravec J., Theurillat J.-P., 2000. International code of the phytosociological nomenclature. 3rd edition. Journal of Vegetation Science 11, 739–768.
- Westhoff V., van der Maarel E., 1973. The Braun-Blanquet approach. 2-nd ed. In: R. Whittaker (Ed.) Classification of Plant Communities, pp. 287–399. Junk, The Hague.
- Winnicki T., 1999. Zbiorowiska roślinne połonin Bieszczadzkiego Parku Narodowego (Bieszczady Zachodnie, Karpaty Wschodnie). Monografie Bieszczadzkie 4: 1–215.